




\_\_\_\_\_ 's  
(your baby's name)

# Immunization Schedule


Print this handy chart to help keep you up-to-date with recommended vaccinations for your little one.



## BIRTH

- First dose Hepatitis B\*

Babies typically receive the first dose of the Hepatitis B vaccine at birth.




## 2 months



- First dose diphtheria, tetanus, and pertussis (Dtap)
- First dose polio (IPV)
- First dose haemophilus influenzae Type B (Hib)
- First dose pneumococcal vaccine (PCV)
- First dose rotavirus
- Second dose Hepatitis B\*

## 4 months

- Third dose Hepatitis B\*
- Second dose diphtheria, tetanus, and pertussis (Dtap)
- Second dose polio (IPV)
- Second dose haemophilus influenzae Type B (Hib)
- Second dose pneumococcal vaccine (PCV)
- Second dose rotavirus



## 6 months



- Third dose diphtheria, tetanus, and pertussis (Dtap)
- Third dose polio (IPV)
- Third dose haemophilus influenzae Type B (Hib)
- Third dose pneumococcal vaccine (PCV)
- Third dose rotavirus
- Flu vaccine\*\*
- Fourth dose Hepatitis B\*

## 12 months

- First dose Hepatitis A
- Measles, mumps, and rubella (MMR)
- Chickenpox (varicella)



## 15 months



- Fourth dose haemophilus influenzae Type B (Hib)
- Fourth dose pneumococcal vaccine (PCV)

Work with your child's pediatrician to find out if he or she missed any necessary vaccines during the first year of life. Contact Children's Hospital of Pittsburgh of UPMC for more information.



\*The CDC only requires three doses of the Hepatitis B immunization, which are typically administered during the first year of a baby's life. Many pediatricians, however, administer four doses when including the Hepatitis B shot as a part of a routine combination vaccine.

\*\*The CDC recommends children age 6 months and older receive an annual flu vaccination. Children receiving the vaccine for the first time are administered a 2-dose series, with each shot separated by 1 month.